



Aalborg Universitet

AALBORG UNIVERSITY
DENMARK

Constraint Counting and Property Predictions in Borophosphate and Phosphosilicate Glasses

Youngman, Randall E.; Hermansen, Christian; Smedskjær, Morten Mattrup; Yue, Yuanzheng

Publication date:
2016

[Link to publication from Aalborg University](#)

Citation for published version (APA):

Youngman, R. E., Hermansen, C., Smedskjær, M. M., & Yue, Y. (2016). *Constraint Counting and Property Predictions in Borophosphate and Phosphosilicate Glasses*. Abstract from 13th International Conference on the Structure of Non-Crystalline Materials, Halifax, Canada.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal -

Take down policy

If you believe that this document breaches copyright please contact us at vbn@aub.aau.dk providing details, and we will remove access to the work immediately and investigate your claim.

Constraint Counting and Property Predictions in Borophosphate and Phosphosilicate Glasses

Randall E. Youngman*, Christian Hermansen, Morten M. Smedskjaer and Yuanzheng Yue

A rapidly growing suite of spectroscopic tools has enabled significant advances in our understanding of short-range structure in glasses. Among these, NMR spectroscopy has been developed to the point where it is routine to identify and quantify different types of network-forming polyhedra, at least for those isotopes possessing favorable NMR properties (e.g. ^{11}B , ^{27}Al , ^{31}P etc...). When used to develop structural models for different glass systems as a function of composition, NMR data on cation speciation is invaluable. Recent work on topological constraint theory in glasses has resulted in more demand for such structural data and models. Atoms in network glasses are constrained by their chemical bonds and bond angles, and the strengths of these constraints depend on the local topology and the chemical nature of the elements. NMR spectroscopic studies of the short-range network structure reveal the identity of these constraints, thus enabling development of a quantitative structural model for borophosphate and phosphosilicate glasses. This combination of detailed structural understanding and topological constraint theory can explain the mixed network former effect (MNFE) in these types of systems, and gives accurate predictions for the composition dependence of glass transition temperature, liquid fragility and indentation hardness.